

*The Interior Highlands Shortleaf Pine Restoration Initiative:
An Overview*



The Interior Highlands Shortleaf Pine Restoration Initiative: An Overview

The Interior Highlands Region:

The Interior Highlands, also referred to here as the Ozark-Ouachita Region, encompasses four broad-scale (Level III) ecoregions, the Ozark Highlands, the Boston Mountains, the Arkansas Valley and the Ouachita Mountains (Omernik, J.M. 1987, U.S. Environmental Protection Agency, 2013; Figure 1).



Figure 1. Ecoregions of the Interior Highlands.

The Ozark Highlands ecoregion occupies most of southern Missouri and extends into northwestern and north-central Arkansas. It is comprised of dissected plateaus that resulted from erosion of a geologic dome uplifted during the Paleozoic era, along with the St. Francis Mountains, the granite core of the Ozarks. The Boston Mountains ecoregion extends from north-central Arkansas into Oklahoma. It is the highest and youngest of the Ozark plateaus, and the most ruggedly dissected. The Ouachita Mountain ecoregion, extending from central Arkansas west into eastern Oklahoma, has been extensively folded and faulted, resulting in the distinct east-west ridges that are evident in the landscape today. The Arkansas Valley ecoregion, a structural valley but much modified by the Arkansas River, lies between the Boston Mountains and Ouachita Mountains in west-central Arkansas, and extends into eastern Oklahoma.

The Ozark-Ouachita Region grades into the Osage Plains to the west and north and into the West Gulf Coastal Plain and Mississippi Alluvial Plain to the south and east. Oak-

hickory forests and woodlands currently predominate throughout much of the Ozarks, with mixed pine-hardwood or pine communities more common in the Ouachitas (Fig. 2). Prairie grasses and forbs are found in the understory of woodlands that are subject to recurrent fire. Glades and barrens occur throughout the region where thin soils and dry exposures limit woody growth. Although glades are characterized by warm-season grasses and a diversity of forbs, eastern red cedar (*Juniperus virginiana*) and Ashe's juniper (*Juniperus ashei*) invade glades during fire-free intervals. Their presence gave rise to the term "cedar glades," which often is used colloquially.

The Ozark Highlands (Level III) ecoregion is comprised of eleven Level IV ecoregions that are differentiated by geologic history, topography, drainage, soils, biota and other features. Elevations range from 144-650 m (400-1800 ft) above sea level, with local relief of 100 or more meters (300 ft) typical of the region. The potential natural vegetation is pine, mixed pine-hardwood, oak-hickory forests and woodlands, prairie and glade-woodland complexes. The Missouri range of short-leaf pine (*Pinus echinata*) is centered in the Ozarks of south-central Missouri (Thom and Wilson 1980).

The Boston Mountains (Level III) ecoregion is comprised of two Level IV ecoregions distinguished by elevation, temperature, precipitation, biota and other features. Gorges and ravines up to 385 m (1250 ft) in depth are common. The juxtaposition of forest and woodland types across the Boston Mountains are determined largely by topography. South-to-west facing slopes on sandstone often are dominated by short-leaf pine mixed with drought tolerant hardwoods such as blackjack oak (*Quercus marilandica*), post oak (*Quercus stellata*), and black hickory (*Carya texana*), while white oak (*Quercus alba*), northern red oak (*Quercus rubra*), and black oak (*Quercus velutina*) are dominant on other slopes. American beech (*Fagus grandifolia*), basswood (*Tilia* sp.), and umbrella magnolia (*Magnolia tripetala*) can be found in deep hollows, ravines, and other mesic sites, principally in the Upper Boston Mountains (Level IV) ecoregion.

The Arkansas Valley ecoregion occupies a transitional zone between the Ozarks and Ouachitas. The (Level III) Arkansas Valley is comprised of four Level IV ecoregions, distinguished by topography, biota and other features. While a large part of the valley is undulating lowland, flat-topped synclinal mountains, remnants of eroded plateaus, and long ridges also are present. The highest elevation in the state is on Magazine Mountain, an isolated, flat-topped mountain within the Arkansas Valley. South-to-west facing slopes on sandstone are often dominated by short-leaf pine along with drought-tolerant hardwoods while upland hardwood forests are dominant on other slopes. Extensive prairies occurred on undulating topography over shale. Bottomland hardwood forests and swamps can be found along the Arkansas, Petit Jean, and Fourche rivers, and seasonally wet-dry hardwood flatwoods occur on poorly-drained flats above the floodplains.

The Ouachita Mountains ecoregion is characteristically a ridge and valley system of east-west trending mountains lying to the south of the Arkansas Valley. The Level III ecoregion is comprised of five Level IV ecoregions distinguished by topography, geographic substrate, drainage, biota and other features. The Fourche Mountains, the most northern of the Level IV ecoregions, occupy more than half of the Natural Division and have the most typical character, with high east-west trending ridges separated by wide valleys having a relatively large, low-gradient stream that follows the structural valley. Local relief can be as much as 540 m (1800 ft). Pine-oak and oak-hickory forests and woodlands occur in repeating east-west bands on the slopes of the ridges, although purer stands of short-leaf pine occur on sites with large areas having few barriers to fire propagation. Prairies appear never to have been common, although bluestem grasses are associated with glades, savannas, and woodlands. Mesic vegetation is restricted to steep, north-facing or otherwise protected slopes.

History of Shortleaf Pine in the Interior Highlands Region:

Shortleaf pine is the only pine native to the Interior Highlands. Shortleaf pine-bluestem and pine-oak woodlands once occurred in very large acreages and across vast landscapes in the Interior Highlands, where the community types were favored by periodic, large-scale fires (Guldin 2007, Guyette et al. 2007). Euro-American settlement of the Ozark/Ouachita region began in the early 1800s, although the population grew more rapidly in the latter part of the century after railroads reached the region (Stroud and Hanson 1981, Smith 1986, Stone County Historical Society 1989, Cunningham 2007). Shortly thereafter, a logging boom ensued, and by 1909, for example, Arkansas ranked 5th in the nation for lumber production (Smith 1986). The bulk of the state's production consisted of short-leaf pine harvested in the Ouachitas. Large volumes of oak were cut in areas where deciduous forests predominated; 500,000 white-oak railroad ties, for example, were shipped via rail from a small town in the White River subdivision of Missouri in 1912 alone (Stone County Historical Society, 1989).

The virgin timber supply was largely exhausted by the 1920s. Natural regeneration occurred, although the stands were characteristically even-aged and/or the original species composition altered. Vast acreages also were planted with pine seedlings. Fire suppression was encouraged throughout the region to protect tree seedlings and saplings (Smith 1986, Palmer 1991). Some cut over areas were converted to agricultural uses, primarily for livestock production in the uplands and to cropland in the bottoms.

Prior to the 1990s, especially on the national forests, even-aged management, or "clear cutting", was the most widely used silvicultural prescription for managing forests; this began to change when the U.S.D.A. Forest Service adopted a "New Perspectives and

Ecosystem Management” approach following a visit by then Chief F. Dale Robertson to the Ouachitas in 1990 (Robertson 2004). Other management techniques (e.g. group selection and shelterwood cuts) began to be tested as methods for managing pine systems in the Interior Highlands during the 1990s, and were the focus of a symposium on ecosystem management in the Interior Highlands in 1999 (Gudlin 2004).

The role of fire in structuring historic shortleaf pine communities also was coming to the attention of managers, but the concept was adopted more slowly. Mark Twain National Forest staff, for example, noticed very positive grass and forb effects following a wildfire in a pine woodlands in the early 1980s. Both excited and curious about the effects of the fire and its implications for the ecosystem, Forest Service staff selected two relatively small sites for pine woodland demonstration areas. They burned a 118 acre site in the spring of 1987, 1989 and 1992 following thinning with horses in 1986, and another 139 acre site in spring of 1988, 1989 and 1992, again following understory thinning (P. Nelson, pers. comm).

The listing of the Red-cockaded Woodpecker, a pine-woodland obligate, as federally endangered played a key role in terms of increasing interest in pineland restoration projects around that time; the development of a recovery plans for the species, beginning in 1985, helped identify the scale at which restoration would be needed, as well as the structural characteristics that managers would need to work toward. The Ouachita National Forest, with a small but extant population of the woodpecker, committed to the restoration of some 250,000 acres of shortleaf pine-bluestem ecosystems with the revisions of their forest plans in 1994, 1996, 2002 and 2005 (Hedrick et al. 2007).

In 1998, ecologists began work to identify sites and landscapes with the best restoration potential in the Missouri Ozarks as well. The Nature Conservancy developed and implemented a rapid ecological assessment technique to identify those sites with the best restoration potential based upon their current condition, remnant vegetation and floristic quality and determined that the only opportunities to recover Missouri’s historic pineries at a large or “landscape” scale were on the Mark Twain National Forest (Ladd et al. 2007). The assessment led to the selection of the Pineknott project area in Carter County, Missouri, initially targeting a tract of more than 12,000 acres, although the thinning and burning needed to restore the woodlands at Pineknott weren’t fully implemented until 2006.

By the early 2000s, pine-bluestem and pine-oak restoration had gained traction among a wide array of conservationists in both Missouri and Arkansas. By 2005, both states had completed their first State Wildlife Action Plans which emphasized natural community restoration of glade-woodland complexes, oak woodlands, and pine and oak-pine woodlands. The number of acres that could be treated, however, was limited by the amount of management agencies on both sides of the state line could fund in addition to

their traditional responsibilities. A cross-border partnership came together and successfully applied for a Doris Duke Foundation grant to not only accelerate the habitat work, but also to foster better regional communication and coordination (Nigh 2007). Once those funds were expended, however, outside funds for restoration work remained largely unavailable and the partnership ceased to meet.

Fortunately, in 2009, an opportunity to secure significant funding for woodland restoration work was made available through passage of Title IV of the Omnibus Public Land Management Act, and establishment of the Collaborative Forest Landscape Restoration Program (CFLRP). The CFLRP, administered by the U.S. Forest Service, was intended to encourage collaborative, science-based ecosystem restoration of priority forest landscapes on and around National Forest lands across the United States. Re-establishing natural fire regimes to reduce the negative consequences of uncharacteristic wildfire also is part of the goal. As a result, up to \$40 million can be appropriated annually from 2009-2019, with up to 4 million a year available for any particular project.

Development of the Interior Highlands SLP Restoration Initiative:

The opportunity to secure significant levels of funding for accelerating shortleaf pine and pine-oak restoration through the Collaborative Forest Landscape Restoration Program, with its emphasis on partnerships and collaboration, provided a renewed incentive for land-managing agencies and organizations with a focus on natural community restoration of shortleaf systems in the Interior Highlands to come together once again. Partnerships formed in the Missouri Ozarks, Arkansas Ozarks, and Ouachitas in support of each national forest's CFLRP proposal, and in January 2011, key federal, state and non-governmental organizations and agencies were brought together by the Central Hardwoods Joint Venture in an effort to communicate and collaborate on shortleaf-related restoration efforts across the Interior Highlands as a whole.¹

A subcommittee consisting of community ecologists from multiple agencies in both states was tasked with developing a set of "desired future conditions" or "DFCs" for pine-bluestem and pine-oak communities as a result of the 2011 meeting. The draft DFCs

¹ The Central Hardwoods Joint Venture, a public-private partnership for bird conservation (see www.chjv.org), has targeted shortleaf pine and pine-oak systems as natural communities capable of providing high-quality habitat for several bird species of conservation concern, including the Red-cockaded Woodpecker, Bachman's Sparrow, Brown-headed Nuthatch, and Prairie Warbler, Northern Bobwhite, and others. While the CHJV's mission is to conserve viable populations of priority bird species within the Central Hardwoods Bird Conservation Region, it also recognizes the importance of shortleaf restoration to support other flora and fauna of conservation concern (see Masters 2007).

were presented to the larger partnership at its second meeting in April 2012 (see Appendix 1). By the time of the April meeting, all three National Forests in the region had been awarded more than \$2 million in funding, via the CFLRP or related programs, for restoration work in 2012, with the potential for that amount or more each year through 2019. If congress continues to support and fund the CFLRP, more than 500,000 acres of pine and pine-oak woodlands should be well on its way toward the restoration goals within the next decade on National Forest lands alone. In more recent years, the partnership has continued to meet periodically to communicate successes and challenges, and consider new needs and opportunities to keep pine restoration moving forward.

The Partnership’s Capacity:

Number of Partners and Varied Expertise: As of spring 2015, numerous state and federal wildlife and land-managing agencies, non-governmental organizations and a university have been represented at meetings of the IHSLPI (Table 1). This diverse group reflects expertise in timber management, restoration ecology, community ecology, research, wildlife conservation and private lands programs, exemplifying the depth and breadth of the partnership and its ability to integrate the multiple facets of a restoration enterprise.

Table 1. Agencies and organizations associated with the Interior Highlands Shortleaf Pine Restoration Initiative.

American Bird Conservancy	Natural Resource Conservation Service
Arkansas Forestry Commission	Oak Woodlands and Forests Fire Consortium
Arkansas Game and Fish Commission	Shortleaf Pine Initiative
Arkansas Natural Heritage Commission	Southeast Conservation Adaptation Strategy
Central Hardwoods Joint Venture	The Nature Conservancy
L-A-D Foundation	University of Missouri, Columbia
Lower Mississippi Valley Joint Venture	U.S. Fish and Wildlife Service
Missouri Department of Conservation	U.S.D.A. Forest Service (National Forests)
Missouri Department of Natural Resources	U.S.D.A. Forest Service (Research Stations)
National Bobwhite Conservation Initiative	

Finally, the vision, mission and operating principles of the IHSLPI have been defined as follows:

Vision

A future where shortleaf pine and associated natural communities regain prominence, enhance biodiversity, and provide sustainable natural resource commodities throughout their historic range within the Interior Highlands region.

Mission

Provide a forum for professional land managers, research scientists and others to work together to advance the restoration of shortleaf pine and pine-oak communities within the Interior Highlands of Arkansas, Missouri and Oklahoma for the benefit of biodiversity and people.

Operating Principles

In working to fulfill our Vision and Mission, the Interior Highlands Shortleaf Pine restoration Initiative embraces the following operating principles:

Science-based Foundation – Our actions and decisions are firmly rooted in science.

Strategic and Adaptive Approach – Restoration is targeted to those sites and landscapes most capable of recovering and sustaining shortleaf pine and associated natural communities over the long term. Our practice of conservation incorporates evaluation and adaptive learning.

Partnerships and Collaboration – Our diverse partnership will share our experiences in shortleaf pine restoration with each other and our constituents to promote effective management practices suitable to the restoration of shortleaf pine and associated natural communities.

Literature cited:

Cunningham, R.J. 2007. Historical and social factors affecting pine management in the Ozark during the late 1800s through 1940. Pgs. 1-7 *In* Kabrick, J.M., D.C. Dey, and D. Gwaze, eds. 2007. Shortleaf pine restoration and ecology in the Ozarks: proceedings of a symposium. Gen. Tech. Rep. NRS-P-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station, 215 p.

Guldin, James M., tech. comp. 2004. Ouachita and Ozarks Mountain symposium: ecosystem management research. Gen. Tech. Rep. SRS-74. Asheville, N.C.: U.S. Department of Agriculture, Forest Service, Southern Research Station, 321. P.

Guldin, James M., 2007. Restoration and management of shortleaf pine in pure and mixed stands – science, empirical observation, and the wishful application of generalities. Pgs. 47-58 *In* Kabrick, J.M., D.C. Dey, and D. Gwaze, eds. 2007. Shortleaf pine restoration and ecology in the Ozarks: proceedings of a symposium. Gen. Tech. Rep. NRS-P-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station, 215 p.

Guyette, R.P., R. Muzika, and S.L. Voelker. 2007. The historical ecology of fire, climate, and the decline of shortleaf pine in the Ozarks. Pgs. 8-18, *In* Kabrick, J.M., D.C. Dey, and D. Gwaze, eds. 2007. Shortleaf pine restoration and ecology in the Ozarks: proceedings of a symposium. Gen. Tech. Rep. NRS-P-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station, 215 p.

Hedrick, L.D., G.A. Bukehofer, W.G. Monatgue, W.F. Pell and J.M. Guldin. 2007. Shortleaf pine-bluestem restoration in the Ouachita National Forest. Pgs. 206-213 *In* Kabrick, J.M., D.C. Dey, and D. Gwaze, eds. 2007. Shortleaf pine restoration and ecology in the Ozarks: proceedings of a symposium. Gen. Tech. Rep. NRS-P-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station, 215 p.

Ladd, D., B. Heumann and P.Nelson. 2007. The pineknot project: restoration of a Missouri Ozark Pinery. Pgs. 204-205 *In* Kabrick, J.M., D.C. Dey, and D. Gwaze, eds. 2007. Shortleaf pine restoration and ecology in the Ozarks: proceedings of a symposium. Gen. Tech. Rep. NRS-P-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station, 215 p.

Masters, R.E. 2007. The importance of shortleaf pine for wildlife and diversity in mixed oak-pine forests and in pine-grassland woodlands. Pgs. 35-46, *In* Kabrick, J.M., D.C. Dey, and D. Gwaze, eds. 2007. Shortleaf pine restoration and ecology in the Ozarks: proceedings of a symposium. Gen. Tech. Rep. NRS-P-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station, 215 p.

McNab, W. H. And P.E. Avers, comps. 1994. Ecological subregions of the United States: Section descriptions. Administrative Publication WO-WSA-5. Washington, DC: U.S. Department of Agriculture, Forest Service. 267 pp.

Nigh, T. A. and W.A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, MO.

Nigh, T.A. 2007. The Ozark Highlands pine-oak woodland restoration partnership. Pgs. 214-215 *In* Kabrick, J.M., D.C. Dey, and D. Gwaze, eds. 2007. Shortleaf pine restoration and ecology in the Ozarks: proceedings of a symposium. Gen. Tech. Rep. NRS-P-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station, 215 p.

Omernik, J.M. 1987. Ecoregions of the conterminous United States. Map (scale 1:7,500,000). *Annals of the Association of American Geographers* 77(1):118-125.

- Palmer, B. 1991. Missouri forests: their history, values and management. Missouri Department of Conservation, Jefferson City, MO.
- Pell, B. 1983. The natural divisions of Arkansas: a revised classification and description. *Natural Areas Journal* 3:12-23.
- Robertson, F. D. 2004. The history of new perspectives and ecosystem management. Pgs. 3-7, *In* Gudlin, James m., tech. comp. 2004. Ouachita and Ozarks Mountain symposium: ecosystem management research. Gen. Tech. Rep. SRS-74. Asheville, N.C.: U.S. Department of Agriculture, Forest Service, Southern Research Station, 321 p.
- Smith, K. L. 1986. Sawmill: the story of the cutting of the last great virgin forest east of the Rockies. Univ. of Arkansas Press. Fayetteville.
- Stone County Historical Society. 1989. History of Stone County Missouri. Stone County Historical Society. Marionville, MO.
- Stroud, H. B. and G. T. Hanson. 1981. Arkansas geography: the physical landscape and the historical-cultural setting. Rose Publishing Co. Little Rock, AR.
- U.S. Environmental Protection Agency. 2013. Level III ecoregions of the continental United States. (map scale 1:7,500,000) U.S. EPA - National Health and Environmental Effects Research Laboratory, Corvallis OR. http://www.epa.gov/wed/pages/ecoregions/level_iii_iv.htm .